

Correlation between Dual-Energy X-Ray Absorptiometry and Panoramic Mandibular Indices in Prediction of Bone Mineral Density in Postmenopausal Females

Safa Hasan Alwan, B.D.S. ⁽¹⁾

Lamia H. Al-Nakib, B.D.S., M.Sc. ⁽²⁾

ABSTRACT

Background: Osteoporosis is a systemic disease of the bone that is characterized by reduced bone mass, which leads to increased bone fragility and fracture particularly in postmenopausal women. The aims of study was to evaluate the relationship between mandibular radiomorphometric indices obtained on digital panoramic radiographs with the bone mineral densities of the lumbar spine evaluated using dual-energy X-ray absorptiometry (DXA) scan, in a population of osteoporotic and non-osteoporotic females.

Materials and methods: In panoramic images obtained from 60 female individuals divided equally into three groups: controls (20-30 years), non-osteoporotic post-menopausal aged 50 years and above and osteoporotic post-menopausal aged 50 years and above, the mean was calculated for mandibular cortical width (MCW), panoramic mandibular index (PMI), mandibular cortical index (MCI) and gonial angle index (GAI) values measured in the right and left sides of the mandible. Bone mineral density (BMD) values were measured by dual energy X-ray absorptiometry (DXA) scan.

Results: Significant positive correlation (r) was observed between bone mineral density of lumbar vertebrae and MCW ($r=0.706$) and PMI ($r=0.668$) of mandible, and a negative correlation was observed between MCI and BMD of lumbar vertebrae ($r=-0.716$). While GAI did not show any significant difference in relation to bone mineral density.

Conclusion: This study showed that MCW, PMI and MCI indices were useful for identifying females with low skeletal (BMD), while GAI was independent in detecting osteoporosis.

Key words: DXA scan, panoramic mandibular indices, post-menopausal females. (J Bagh Coll Dentistry 2016; 28(3):87-91).

INTRODUCTION

Osteoporosis is a disease characterized by low bone mass and micro-architectural deterioration of bone tissue, leading to bone fragility and enhanced susceptibility to fractures. Osteoporosis is predominantly a condition of the elderly⁽¹⁾. It affects mostly women, especially after menopause as a result of estrogen withdrawal⁽²⁾. A major obstacle to combating osteoporosis is the failure to identify individuals who have osteoporosis until the clinical consequences of osteoporosis have occurred (i.e., fractures form with little trauma to the bones)⁽³⁾.

BMD evaluation by dual-energy X-ray absorptiometry (DXA) testing is considered the gold standard for fracture risk prediction⁽⁴⁾. However, in addition to not being recommended by the WHO as a triage screening tool for osteoporosis, it has a high financial cost⁽⁵⁾. Because the bones of the oral cavity are similar in structure and physiology to various other bones in the skeleton, several studies have been conducted with the object of detecting whether these skeletal changes in the mandible are specific to the osteoporotic stage^(4,5).

Panoramic radiography has been an important component of dental diagnostic radiology for over 40 years⁽⁶⁾.

Recent clinical studies have shown that panoramic radiography plays a critical role in the identification and evaluation of osteoporotic patients or those with low BMD by dentists⁽⁷⁾.

Qualitative and quantitative indices which include the mandibular cortical index (MCI), mandibular cortical thickness (MCW), gonial angle index (GAI) or panoramic mandibular index (PMI) have been used for panoramic radiographs, to assess the bone quality and to observe signs of resorption and osteoporosis⁽⁶⁾.

The present study focuses on obtaining a possible mean of early detection of osteoporosis by panoramic radiography.

MATERIALS AND METHODS

A cross-sectional study was conducted on 60 Iraqi females who had been referred to bone densitometer center for BMD evaluation by DXA technique at the X-Ray Institute in the Medical City in Baghdad.

The patients were asked to participate in this clinical trial as volunteers, and they were thoroughly informed about the procedure. The sample was divided into three groups according to their age and osteoporotic status:

- **1st group:** 20 females with age 20-30 as control group (non-osteoporotic).
- **2nd group:** 20 postmenopausal non-osteoporotic females with age from 50 and above.

(1) M.Sc. student, Department of Oral Diagnosis, College of Dentistry, University of Baghdad.

(2) Assistant professor, Department of Oral Diagnosis, College of Dentistry, University of Baghdad.

- **3rd group:** 20 postmenopausal osteoporotic females with age from 50 and above.

Females with natural menopause (which occurs after 12 months of amenorrhoea and for which there was no obvious pathologic cause) with no history of hysterectomy, were included in the study. Smoking, alcoholism and patients with any known systemic disease that would affect bone metabolism like hyperparathyroidism, hypoparathyroidism, hyperthyroidism, Paget's disease, osteomalacia, renal osteodystrophy, cancers with bone metastasis or significant renal impairment and patients who were on specific drugs (corticosteroids) which are known to have adverse effects on bone metabolism were excluded from the study.

Dual-energy X-ray absorptiometry (DEXA) scan of the spinal vertebrae (L2-L4) was performed by using osteosys DEXXUM 3 (Korea) machine. This procedure is the current gold standard for measuring bone mass and detecting osteoporosis⁽⁸⁾.

Thereafter, The patients were then subjected to panoramic imaging in the Department of Oral Medicine, Diagnosis and Radiology at the College of Dentistry (Baghdad University), on Dimax 3 Digital X-ray machine manufactured by Planmeca Oy, Helsinki, Finland. After that, the images were manipulated on the computer monitor of the x-ray machine to achieve best image quality, then converted to **JPG** (joint photographic experts group) files; so that linear and angular measurements were calculated by AutoCAD software (2007).

The results of DXA scan were then compared with the results of the indices from the panoramic radiographs.

The following radiomorphometric indices were measured for each patient:

- **Mandibular cortical width (MCW)**, which is the measurement of the cortical width at the mental foramen region⁽⁹⁾.
- **Panoramic mandibular index (PMI)** is the ratio of the thickness of the mandibular cortex to the distance between the superior margin of mental foramen and the inferior mandibular cortex, presented by Benson *et al.*⁽¹⁰⁾.
- **Mandibular cortical index (MCI)** refers to the mandibular cortical shapes on dental

panoramic radiographs, and is categorized into one of the three groups according to the method of Klemetti *et al.*, as follows: C1: Normal cortex, C2: Mild to moderately eroded cortex and C3: severely eroded cortex⁽¹¹⁾.

- **Gonial angle index (GAI)**, which refers to the size of the gonial angle of mandible⁽¹²⁾.

The WHO diagnostic criteria for osteoporosis defines osteoporosis in terms of a T-score which is below -2.5 and osteopaenia in terms of a T-score which is between -2.5 and -1⁽¹³⁾.

RESULTS

The results showed that MCW and PMI mean value was the highest for the healthy young female group (group I). While, the MCW and PMI value of post-menopausal osteoporotic female group (group III) showed the lowest mean. The results showed that all the three groups were statistically highly significant in MCW and PMI as P-value < 0.001.

The mean GAI value for group I was the lowest among the rest of the study groups. While the GAI of group III showed the highest value. The results showed that all the three groups were statistically non-significant in GAI as p-value was 0.21 (Table 1).

Regarding MCI, it was found that group I has normal mandibular cortical index (C1). While group II shows lacunar resorption (C2) in 15% of the cases. And group III shows lacunar resorption (C2) in 45% and clear porosity (C3) in 40% of the cases. The difference between the 3 groups showed a high statistical significance as the p-value < 0.001 (Table 2).

The results showed a positive correlation between MCW, PMI and T-score as r pearson value was (0.706) and (0.668), respectively. And a strong negative correlation between MCI and T-score as r value was (-0.716). There is a weak negative correlation between GAI and T-score as r value (-0.224); (Table 3).

DISCUSSION

Panoramic radiography is a routine imaging method in dentistry and is part of many recall programmes.

Table1: Mean values of Mandibular cortical width, Panoramic mandibular index and Gonial angle index according to DEXA results.

Groups	MCW			PMI			GAI		
	Mean	SD	P-value	Mean	SD	P-value	Mean	SD	P-value
Normal young group	3.8	0.38	<0.001	0.246	0.032	<0.001	121.9	7.34	0.21 [NS]
Post-menopausal normal group	3.4	0.43		0.207	0.032		123.6	6.74	
Post-menopausal osteoporotic group	2.3	0.61		0.141	0.043		126	7.97	

Table2: Showing frequency and percentage of MCI in different study groups.

	Healthy young group		Postmenopausal non Osteoporotic group		Postmenopausal osteoporotic group	
	N	%	N	%	N	%
MCI						
C1	20	100.0	17	85.0	3	15.0
C2	0	0.0	3	15.0	9	45.0
C3	0	0.0	0	0.0	8	40.0
Total	20	100.0	20	100.0	20	100.0

Table3: Linear correlation coefficient of MCI, PMI, GAI, T-score and age of all three groups

	MCW	PMI	GAI	MCI
T-score	r=0.706 P<0.001	r=0.668 P<0.001	r=-0.224 P=0.17	r=-0.716 P<0.001

Several studies suggest that panoramic radiomorphometric indices may be useful for identifying patients with low skeletal bone mineral density or osteoporosis^(14,15). Although some studies have found no relationship between skeletal and mandibular BMD^(7,16).

Measurement of the thickness of the mandibular cortical width in panoramic radiographs has been suggested as a way to predict patients with low bone mineral density⁽¹⁷⁾.

In the current study, the cortical bone in the mental region was significantly thinner in post-menopausal osteoporotic females if compared with post-menopausal non-osteoporotic females. This result was expected as it concurs with previous studies⁽¹⁸⁻²⁰⁾.

Besides that, MCW was also significantly thinner in post-menopausal groups than the young healthy group, and this can be attributed to the fact that MCW is affected by hormonal changes like estrogens which have an important role in the regulation of skeletal development and homeostasis; this is demonstrated by the dramatic loss of bone that occurs after menopause. (During estrogen deficiency there is prolongation of osteoclast life span due to inhibition of apoptosis)^(21,22). This result was in agreement with Taguchi *et al.* and Khojastehpour *et al.*, who stated that age was shown to have a significant correlation with the MCW, as age increased, there was a decreasing rate in cortical width^(15,23).

Panoramic mandibular index was first proposed by Benson *et al.*, as radiomorphometric index of adult cortical bone mass⁽¹⁰⁾. Predictably, in this study PMI demonstrated similar age-related

correlations as those of MCW by showing a statistically significant difference among all the three groups; the result of this study agrees with that of Halling *et al.* and Kim *et al.*^(24,25).

The results of this study is consistent with the results of Horner and Devlin, in which the PMI values measured in females with osteoporosis on panoramic radiographs were compared with the mandibular bone values measured with DXA, and a significant relationship was found between the two. They concluded that PMI could be used as an indicator of mandibular bone density⁽¹⁸⁾. Our results disagrees with the study done by Drozdowska *et al.*, which showed that there was no correlation between PMI and DEXA measurement and they suggested that it should not be used as an indicator of skeletal status⁽⁷⁾.

Regarding the gonial angle, in the present study, the size of the gonial angle increases as age increases, but it did not show a statistical significant difference. Which agrees with the study conducted by Dutra *et al.*, done on British population, where the correlation between age and gonial angle was statistically non-significant⁽²⁶⁾, also with Ceylan *et al.*, where the gonial angle did not increase as age increases in the edentulous individuals; therefore, they stated that lack of correlation between the gonial angle and the age indicates that the angle does not change with increase of the individual's age⁽²⁷⁾.

The present study disagrees with that done by Mahdi and Al-Nakib, which showed that older subjects had significantly larger gonial angle than younger ones⁽²⁸⁾.

One of the most commonly studied parameters of mandibular bone with respect to osteoporosis is the porosity of the mandibular cortical bone. In the present study, C1 shape of cortex was seen in younger females, but as age increased, the number of individuals who had C2 and C3 categories increased, presumably reflecting age related bone loss.

In the present study, C2 and C3 categories were predominantly seen in post-menopausal osteoporotic females, which were again supported by previous studies⁽²⁹⁻³¹⁾.

Klemetti *et al.* evaluated the MCI, which was also known by the author's name. Our results confirm the results achieved by Klemetti *et al.*, as they suggested that a thin or eroded inferior cortex of the mandible detected on dental panoramic radiographs, an indicator of alterations of the mandible, is useful for identifying post-menopausal females with undetected low skeletal BMD or osteoporosis⁽¹¹⁾.

Regarding T-score, we found that there is a positive correlation between MCW and T-score, this correlation was statistically highly significant, and this agrees with Hekmatin *et al.*, who found a significant correlation between BMD and MCW and positive correlation between MCW and T-score⁽³²⁾. T-score was also correlated positively with the PMI, which statistically has highly significant difference that was repeated by Parlani⁽³³⁾. T-score, in our study also correlates positively with the MCI, and this is agreed by Gulsahiet *al.*⁽³⁰⁾.

From this study, we can conclude that panoramic radiography gives sufficient information to make an early diagnosis regarding osteoporosis in post-menopausal females, and this may be helpful in the prevention of osteoporotic fractures in elderly females.

REFERENCES

- Rizzoli R, Bruyere O, Cannata-Andia JB, Devogelaer JP, Lyritis G, Ringe JD, Vellas B, Reginster JY. Management of osteoporosis in the elderly. *Curr Med Res Opin* 2009;25: 2373-87.
- Karasik D. Osteoporosis: an evolutionary perspective. *Hum Genet* 2008;124: 349-56.
- NIH Consensus Development Panel on Osteoporosis Prevention, Diagnosis, and Therapy *JAMA* 2001;285: 785-95.
- Horner K, Karayianni K, Mitsea A, Berkas L, Mastoris M, Jacobs R, et al. The mandibular cortex on radiographs as a tool for osteoporosis risk assessment: the osteodent Project. *J Clin Densitom* 2007; 10: 138-146.
- Taguchi A. Triage screening for osteoporosis in dental clinics using panoramic radiographs. *Oral Dis* 2010; 16: 316-27.
- Yasar F, Senar S, Yesilova E, Akgunlu F. Mandibular cortical index evaluation in masked and unmasked panoramic radiographs. *Dentomaxillofac Radiol* 2009;38:86-91.
- Drozdowska B, Pluskiewicz W, Tarnawska B., Panoramic-base mandibular indices in relation to mandibular bone mineral density and skeletal status assessed by dual energy X-ray absorptiometry and quantitative ultrasound. *DMFR* 2002; 31: 361-7.
- Leite AF, Figueiredo PT, Guia CM, Melo NS, de Paula AP. Correlations between seven panoramic radiomorphometric indices and bone mineral density in post-menopausal women. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2010; 109: 449-56.
- Ledgerton D, Horner K, Devlin H, Worthington H. Panoramic mandibular index as a radiomorphometric tool: an assessment of precision. *Dentomaxillofac Radiol* 1997;26:95-100.
- Benson BW, Prihoda TJ, Glass BJ. Variations in adult cortical bone mass as measured by a panoramic mandibular index. *Oral Surg Oral Med Oral Pathol* 1991;71: 349-56.
- Klemetti E, Kolmakov S, Kroeger H. Pantomography in assessment of the osteoporosis risk group. *Scand J Dent Res* 1994; 102: 68-72.
- Ceiiareginawinckmahl Comparison of morphometric indices obtained from dental panoramic radiography of identifying individuals with osteoporosis or osteopenia. *Radiolbris* 2008;41:3.
- El Maghraoui A, Roux C. DXA scanning in clinical practice. *QJM* 2008; 101:605-17.
- Bollen AM, Taguchi A, Hujuel PP, Hollender LG. Case-control study on self-reported osteoporotic fractures and mandibular cortical bone. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2000; 90: 518-24.
- Taguchi A, Suei Y, Ohtsuka M, Otani K, Tan-imoto K, Ohtaki M. Usefulness of panoramic radiography in the diagnosis of postmenopausal osteoporosis in women. Width and morphology of inferior cortex of the mandible. *Dentomaxillofac Radiol* 1996; 25: 263-7.
- Devlin CV, Horner K, Devlin H. Variability in measurement of radiomorphometric indices by general dental practitioners. *Dentomaxillofac Radiol* 2001; 30: 120-5.
- Dutra V, Yang J, Devlin H, Susin C. Radiomorphometric indices and their relation to gender, age, and dental status. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2005;99:479-84.
- Horner K, Devlin H. The relationship between mandibular bone mineral density and panoramic radiographic measurements. *J Dent* 1998;26: 337-43.
- Taguchi A, Tsuda M, Ohtsuka M, Kodama I, Sanada M, Nakamoto T, Inagaki K, Noguchi T, Kudo Y, Suei Y, Tanimoto K, Bollen AM. Use of dental panoramic radiographs in identifying younger postmenopausal women with osteoporosis. *Osteoporos Int* 2006;17: 387-94.
- Bodade PR, Mody RN. Panoramic radiography for screening postmenopausal osteoporosis in india: A Pilot Study. *OHDM* 2013;12:72-65.
- Haslett C, Braum WE, Randolph T, et al. (Davidson's): Principles and Practice of Medicine. 19thed. London: Hunter; 2005. pp. 1025-7.
- Marcus R, Feldman D, Nelson DA, Rosen CJ. Osteoporosis. 3rded. Elsevier; 2008. pp. 3,1031,1416-1441.
- Khojastehpour L, Afssa M, Dabbaghmanesh MH. Evaluation of Correlation between Width and

- Morphology of Mandibular Inferior Cortex in Digital Panoramic Radiography and Postmenopausal Osteoporosis. *Iranian Red Crescent Medical J* 2011;13(3):181-6.
24. Halling A, Perrson GR, Berglund J, Johansson O, Renvert S. Comparison between the Klemetti index and heel DXA BMD measurements in the diagnosis of reduced skeletal bone mineral density in the elderly. *Osteoporosis Int* 2005;16:999-1003.
 25. Kim Y, Kim K, Koh K. The relationship between age and the mandibular cortical bone thickness by using panoramic radiograph. *Korean J Oral Maxillofac Radiol* 2010; 40:83-7.
 26. Dutra V, Devlin H, Susin C, Yang J, Horner K., Mandibular morphological changes in low bone mass edentulous females: evaluation of panoramic radiographs. *Oral Surg Oral Med Oral Pathol Oral Radiol Endodontology* 2006; 102:663-8.
 27. Ceylan C, Yanikoglu N, Yilmaz A, Ceylan Y, .Changes in the mandibular angle in the dentulous and edentulous states. *J Prosthet Dent* 1998; 80: 680-4.
 28. Mahdi AS, Al-Nakib LH. Assessment of mandibular radiomorphometric indices as predictors of osteoporosis in postmenopausal women (Cephalometric reconstructed computed tomographical study). *JBCD* 2014; 26(2): 50-7.
 29. Geraels WGM, Verheij JGC, van der Stelt PF, Horner K, Lindh C, Nicopoulou-Karayianni K, et al. Osteoporosis and the general dental practitioner: reliability of some digital dental radiological measures. *Community Dent Oral Epidemiol* 2007;35:465-71.
 30. Gulsahi A, Yuzugullu B, Imirzahoglu P, Gene Y. Assessment of panoramic radiomorphometric indices in Turkish patients of different age groups, gender and dental status. *Dentomaxillofac Radiol* 2008;37:288-92.
 31. Dagistan S, Bilge OM. Comparison of antegonial index, mental index, panoramic mandibular index and mandibular cortical index values in the panoramic radiographs of normal males and male patients with osteoporosis. *Dentomaxillofac Radiol* 2010;39: 290-4.
 32. Hekmatin E, Ahmadi SS, Ataiekhorsgani M, Feizianfard M, Jafaripozve S, Jafaripozve N., Prediction of lumbar spine bone mineral density from the mandibular cortical width in postmenopausal women. *J Res Med Sci* 2013; 18: 951-5.
 33. Parlani S, Nair P, Agrawal S, Chitumalla R, Beohar G, Katar U. Role of panoramic radiographs in the detection of osteoporosis. *Oral Hyg Health IJGM* 2014; 2: 2332-0672